

Comment on “Universal relation between the Kolmogorov-Sinai Entropy and the thermodynamic entropy in simple liquids”

In an intriguing paper [1], Dzугutov, Aurell and Vulpiani (DAV) establish a quantitative connection in the form of a universal linear law between the Kolmogorov-Sinai entropy h_{KS} and the excess entropy S_{ex} for three spherical potentials: $h_{KS}^{DAV}/\Gamma_{coll} = 0.62 + 1.106S_{ex}$ (1). Here Γ_{coll} is the average pair collision frequency of an equivalent hard sphere fluid. We applied this relation to a hard sphere fluid, comparing it with the values of h_{KS}^{DP} given by Dellago and Posch (DP) [2]. In Fig. 1 we plot h_{KS}^{DP} , as well as (1). In the calculation of the curve labelled DAV we used $S_{ex} = 3 - 12(9 - \pi n\sigma^3)/(6 - \pi n\sigma^3)^2$ based on the Carnahan-Starling formula for the pressure of a hard sphere fluid and on thermodynamic integration [1]. It is clear that (1) does not apply to the h_{KS}^{DP} . The same holds for the case of hard disk systems in two dimensions, for which the assumption of any linear law like (1), leads to the same contradictions.




FIG. 1: Reduced h_{KS} values for hard sphere fluids, labelled as in the text. Here m is the mass of the particles, n their number density, σ their radius, N their number and K their kinetic energy. The range of $n\sigma^3$ used by DAV is indicated by the vertical lines.

Paper [1] was inspired by an earlier one of Dzугutov [3] concerning a universal connection between the self diffusion coefficient D and S_{ex} of a number of dense (including hard sphere) fluids, of the form: $D^* \equiv D/\Gamma\sigma^2 = 0.049e^{S_{ex}}$ (2), where σ is an effective hard sphere diameter. The range of density considered in [2] is $0.4 \leq n\sigma^3 \leq 0.9$, i.e. restricted to high fluid densities, but as shown in Fig.2 the agreement of (2) and the numerical data from Alder, Gass and Wainwright (AGW) [4] and Erpenbeck and Wood (EW) [5] for hard spheres is not good. The problem is shown in a wider range in Fig. 3 where (2), transformed to D/D_B , is compared with the numerical data of [4,5]. Here D_B is the Boltzmann value for the self diffusion coefficient. Note that D/D_B does not monotonically decrease with density, which is unphysical. Never-

theless, a connection between dynamical (h_{KS} , D) and thermodynamical (S_{ex}) quantities remains a challenge.

The authors would like to thank I.M. de Schepper for very helpful correspondence.




FIG. 2: Comparison of D^* for hard spheres from the numerical data of AGW and EW with those from (2) (Dz.), in the range of S_{ex} as considered by Dzугutov. Here the real S_{ex} values are used, instead of the approximation of Dzугutov.




FIG. 3: Comparison between the numerical data of AGW and EW for D/D_B for hard spheres and those obtained from (2). The vertical lines indicate the range considered by Dzугutov.

E.G.D. Cohen, Rockefeller University, New York, NY 10021 - U.S.A.

L. Rondoni, Politecnico di Torino, C. D. Abruzzi 24, I-10129 Torino - Italy

PACS numbers: 05.70Ce, 05.45.+b, 61.20.Ja

- [1] M. Dzугutov, et al. Phys. Rev. Lett. **81**, 1762 (1998)
- [2] Ch. Dellago and H.A. Posch, Physica A, **240**, 68 (1997)
- [3] M. Dzугutov, Nature, **381**, 137 (1996)
- [4] B.J. Alder, et al. J. Chem. Phys. **53**, 3813 (1970)
- [5] J.J. Erpenbeck et al. Phys. Rev. A **43**, 4254 (1991)